


Amendments to the Claims:

The listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Currently Amended) A substituted 4-aminocyclohexanol compound corresponding to formula I,



wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl; wherein R¹ and R² are not both H, or the radicals R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁵CH₂CH₂ or (CH₂)₃₋₆,

wherein

R⁵ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl;

R³ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;
R⁴ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl; -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

wherein

Y = O, S or H₂;

R⁶ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₇-alkyl; or saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₆-alkyl;

R⁷ represents H; unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl;

R⁸ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;

L represents -C(O)-NH-, -NH-C(O)-, -C(O)-O-, -O-C(O)-, -O-, -S- or -S(O)₂-; and

R⁹ represents unsubstituted or singly or multiply substituted aryl or heteroaryl,

provided that,

~~if R3 = substituted or unsubstituted phenyl, and R4 = phenyl or -CHR6R7, -CHR6-CH2R7, -CHR6-CH2-CH2R7, -CHR6-CH2-CH2-CH2R7, -C(Y)R7, -C(Y)-CH2R7, -C(Y)-CH2-CH2R7 or -C(Y)-CH2-CH2-CH2R7~~

~~where Y = H₂,~~

~~R6 = H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C1-5-alkyl,~~

~~or~~

~~R7 = H, substituted or unsubstituted C3-8 cycloalkyl or phenyl,~~

~~R1 and R2 independently of one another are not both C1-5 alkyl, R1 and R2 independently of one another are not both C1-5 alkyl,~~

~~or~~

~~if R3 = substituted or unsubstituted thiophenyl, and R4 = -CH2-CH2-phenyl,~~

~~the radicals R1 and R2 do not together form a ring and represent (CH2)5,~~

~~or a salt thereof with a physiologically tolerated acid.~~

or an acid, base, solvate or a physiologically acceptable salt thereof

provided that,

if R³ = substituted or unsubstituted phenyl, and R⁴ = phenyl or -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷

where Y = H₂,

R⁶ = H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₅-alkyl,

or

R⁷ = H, substituted or unsubstituted C₃₋₈-cycloalkyl or phenyl, then


R¹ and R² independently of one another are not both C₁₋₅-alkyl,

or

if R³ = substituted or unsubstituted thiophenyl, and R⁴ = -CH₂-CH₂-phenyl, then

the radicals R¹ and R² do not together form a ring and represent (CH₂)₅.

2. (Previously Presented) A substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1,



wherein

the radicals R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁵CH₂CH₂ or (CH₂)₃₋₆,

where R⁵ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl;

R³ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;

R⁴ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl; -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

wherein

Y = O, S or H₂;

R⁶ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₇-alkyl; or saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₆-alkyl;

R⁷ represents H; unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl;

R⁸ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;


L represents -C(O)-NH-, -NH-C(O)-, -C(O)-O-, -O-C(O)-, -O-, -S- or -S(O)₂-; and

R⁹ represents unsubstituted or singly or multiply substituted aryl or heteroaryl,

provided that,

if R³ = substituted or unsubstituted thiophenyl, and R⁴ = -CH₂-CH₂-phenyl, the radicals R¹ and R² do not together form a ring and represent (CH₂)₅, or a salt thereof with a physiologically tolerated acid.

3. (Currently Amended) A substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1,



I

wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl; wherein R¹ and R² are not both H,

R³ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;

R⁴ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl

or heteroaryl; -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

wherein

Y = O, S or H₂;

R⁶ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₇-alkyl; or saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₆-alkyl;

R⁷ represents H; unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl;

R⁸ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;

L represents -C(O)-NH-, -NH-C(O)-, -C(O)-O-, -O-C(O)-, -O-, -S- or -S(O)₂-; and

R⁹ represents unsubstituted or singly or multiply substituted aryl or heteroaryl,

provided that,

~~if R3 = substituted or unsubstituted phenyl, and R4 = phenyl or -CHR₆R₇, -CHR₆-CH₂R₇, -CHR₆-CH₂-CH₂R₇, -CHR₆-CH₂-CH₂-CH₂R₇, -C(Y)R₇, -C(Y)-CH₂R₇ or -C(Y)-CH₂-CH₂R₇~~

~~where Y = H₂,~~

~~R₆ = H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₅ alkyl,~~

~~or~~

~~R₇ = H, substituted or unsubstituted C₃₋₈ cycloalkyl or phenyl,~~

~~R₁ and R₂ independently of one another are not both C₁₋₅ alkyl, R₁ and R₂~~

~~independently of one another are not both C₁₋₅-alkyl,~~

~~or~~

~~if R³ = substituted or unsubstituted thiophenyl, and R⁴ = -CH₂CH₂-phenyl,
the radicals R¹ and R² do not together form a ring and represent (CH₂)₅,
or a salt thereof with a physiologically tolerated acid.~~

or an acid, base, solvate or a physiologically acceptable salt thereof
provided that,

if R³ = substituted or unsubstituted phenyl, and R⁴ = phenyl or -CHR⁶R⁷, -
CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-
CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷

where Y = H₂,

R⁶ = H, saturated or unsaturated, branched or unbranched, singly
or multiply substituted or unsubstituted C₁₋₅-alkyl,

or


R⁷ = H, substituted or unsubstituted C₃₋₈-cycloalkyl or phenyl, then
R¹ and R² independently of one another are not both C₁₋₅-alkyl,

or

if R³ = substituted or unsubstituted thiophenyl, and R⁴ = -CH₂-CH₂-
phenyl, then

the radicals R¹ and R² do not together form a ring and represent (CH₂)₅.

4. (Previously Presented) A substituted 4-aminocyclohexanol corresponding to formula I of claim 1,



I

wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl; wherein R¹ and R² are not both H, or the radicals R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁵CH₂CH₂ or (CH₂)₃₋₆,

wherein R⁵ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl;

R³ represents unsubstituted or singly or multiply substituted heteroaryl;

R⁴ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl; -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

wherein

Y = O, S or H₂;

R⁶ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₇-alkyl; or saturated or

unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₆-alkyl;

R⁷ represents H; unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl;

R⁸ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;


L represents -C(O)-NH-, -NH-C(O)-, -C(O)-O-, -O-C(O)-, -O-, -S- or -S(O)₂-; and

R⁹ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;

provided that,

if R³ = substituted or unsubstituted thiophenyl, and R⁴ = -CH₂-CH₂-phenyl, the radicals R¹ and R² do not together form a ring and represent (CH₂)₅, or a salt thereof with a physiologically tolerated acid.

5. (Currently Amended) A substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1,



wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or

heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl; wherein R¹ and R² are not both H, or the radicals R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁵CH₂CH₂ or (CH₂)₃₋₆,

wherein R⁵ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl;

R³ represents unsubstituted or singly or multiply substituted aryl;

R⁴ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl; -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

wherein

Y = O, S or H₂;

R⁶ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₇-alkyl; or saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₆-alkyl;

R⁷ represents H; unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl;

R⁸ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;

L represents -C(O)-NH-, -NH-C(O)-, -C(O)-O-, -O-C(O)-, -O-, -S- or -S(O)₂-; and

R⁹ represents unsubstituted or singly or multiply substituted aryl or

heteroaryl,

provided that,

~~if R₃ = substituted or unsubstituted phenyl, and R₄ = phenyl or -CHR₆R₇, -CHR₆-CH₂R₇, -CHR₆-CH₂-CH₂R₇, -CHR₆-CH₂-CH₂-CH₂R₇, -C(Y)R₇, -C(Y)-CH₂R₇, -C(Y)-CH₂-CH₂R₇ or -C(Y)-CH₂-CH₂-CH₂R₇~~

~~where Y = H₂,~~

~~R₆ = H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₅ alkyl,~~

~~or~~

~~R₇ = H, substituted or unsubstituted C₃₋₈-cycloalkyl or phenyl,~~

~~R₁ and R₂ independently of one another are not both C₁₋₅ alkyl, R₁ and R₂ independently of one another are not both C₁₋₅ alkyl,~~

~~or~~

~~if R₃ = substituted or unsubstituted thiophenyl, and R₄ = -CH₂-CH₂ phenyl,~~

~~the radicals R₁ and R₂ do not together form a ring and represent (CH₂)₅,~~

~~or a salt thereof with a physiologically tolerated acid.~~

or an acid, base, solvate or a physiologically acceptable salt thereof

provided that,

if R³ = substituted or unsubstituted phenyl, and R⁴ = phenyl or -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷

where Y = H₂,

R⁶ = H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₅-alkyl,

or

R⁷ = H, substituted or unsubstituted C₃₋₈-cycloalkyl or phenyl, then


R¹ and R² independently of one another are not both C₁₋₅-alkyl,

or

if R³ = substituted or unsubstituted thiophenyl, and R⁴ = -CH₂-CH₂-phenyl, then

the radicals R¹ and R² do not together form a ring and represent (CH₂)₅.

6. (Previously Presented) A substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1,



wherein

the radicals R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁵CH₂CH₂ or (CH₂)₃₋₆,

wherein R⁵ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl;

R³ represents unsubstituted or singly or multiply substituted aryl;

R⁴ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl; -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

wherein

Y = O, S or H₂;

R⁶ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₇-alkyl; or saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₆-alkyl;


R⁷ represents H; unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl;

R⁸ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;

L represents -C(O)-NH-, -NH-C(O)-, -C(O)-O-, -O-C(O)-, -O-, -S- or -S(O)₂-; and

R⁹ unsubstituted or singly or multiply substituted aryl or heteroaryl, or a salt thereof with a physiologically tolerated acid.

7. (Previously Presented) A substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1,



wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or

heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl; wherein R¹ and R² are not both H, R³ represents unsubstituted or singly or multiply substituted heteroaryl; R⁴ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl; -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

wherein

Y = O, S or H₂;

R⁶ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₇-alkyl; or saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₆-alkyl;

R⁷ represents H; unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl;


R⁸ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;

L represents -C(O)-NH-, -NH-C(O)-, -C(O)-O-, -O-C(O)-, -O-, -S- or -S(O)₂-; and

R⁹ represents unsubstituted or singly or multiply substituted aryl or heteroaryl,

or a salt thereof with a physiologically tolerated acid.

8. (Previously Presented) A substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1,



I

wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl; wherein R¹ and R² are not both H, or the radicals R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁵CH₂CH₂ or (CH₂)₃₋₆,

wherein R⁵ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl;

R³ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;

R⁴ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl; -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

wherein

Y = H₂,

R⁶ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₇-alkyl

R⁷ represents unsubstituted or singly or multiply substituted heteroaryl,


R⁸ represents unsubstituted or singly or multiply substituted aryl or heteroaryl,

L represents -C(O)-NH-, -NH-C(O)-, -C(O)-O-, -O-C(O)-, -O-, -S- or -S(O)₂-; and

R⁹ represents unsubstituted or singly or multiply substituted aryl or heteroaryl,

or a salt thereof with a physiologically tolerated acid.

9. (Previously Presented) A substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1,



wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl; wherein R¹ and R² are not both H,

or the radicals R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁵CH₂CH₂ or (CH₂)₃₋₆;

wherein R⁵ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or

C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl;

R³ represents unsubstituted or singly or multiply substituted aryl or heteroaryl;


R⁴ represents -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷ or -CHR⁶-CH₂-CH₂-CH₂R⁷;

R⁶ represents saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₆-alkyl; and

R⁷ represents H; respectively unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl,

or a salt thereof with a physiologically tolerated acid.

10. (Previously Presented) A substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1,



wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl; wherein R¹ and R² are not both H, or the radicals R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂,

$\text{CH}_2\text{CH}_2\text{NR}^5\text{CH}_2\text{CH}_2$ or $(\text{CH}_2)_{3-6}$,

wherein R^5 represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C_{1-8} -alkyl or C_{3-8} -cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C_{1-3} -alkylene, C_{3-8} -cycloalkyl or heteroaryl;

R^3 represents unsubstituted or singly or multiply substituted aryl or heteroaryl;

R^4 represents $-\text{C}(\text{Y})\text{R}^7$, $-\text{C}(\text{Y})\text{-CH}_2\text{R}^7$, $-\text{C}(\text{Y})\text{-CH}_2\text{-CH}_2\text{R}^7$ or $-\text{C}(\text{Y})\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{R}^7$;

wherein

$\text{Y} = \text{O}$ or S ; and

R^7 represents H; unsubstituted or singly or multiply substituted C_{3-8} -cycloalkyl, aryl or heteroaryl,

or a salt thereof with a physiologically tolerated acid.

11. (Original) A substituted 4-aminocyclohexanol compound according to claim 1 wherein:

R^1 and R^2 independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C_{1-8} -alkyl; wherein R^1 and R^2 are not both H,

or the radicals R^1 and R^2 together form a ring and represent $\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2$, $\text{CH}_2\text{CH}_2\text{NR}^5\text{CH}_2\text{CH}_2$ or $(\text{CH}_2)_{3-6}$,

where R^5 represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C_{1-8} -alkyl.

12. – 124. (Cancelled)

125. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 3, wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl; wherein R¹ and R² are not both H.

126. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 7, wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl; wherein R¹ and R² are not both H.

127. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 2, wherein

R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁵CH₂CH₂ or (CH₂)₃₋₆,

wherein R⁵ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl.

128. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 6, wherein

R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁵CH₂CH₂ or (CH₂)₃₋₆,

wherein R⁵ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl.

129. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 1, wherein

R^3 represents unsubstituted or singly or multiply substituted phenyl, naphthyl, anthracenyl, thiophenyl, benzothiophenyl, pyridyl, furyl, benzofuranyl, benzodioxolanyl, indolyl, indanyl, benzodioxanyl, pyrrolyl, pyrimidyl or pyrazinyl.

130. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 4, wherein

R^3 represents unsubstituted or singly or multiply substituted thiophenyl, benzothiophenyl, pyridyl, furyl, benzofuranyl, benzodioxolanyl, indolyl, indanyl, benzodioxanyl, pyrrolyl, pyrimidyl or pyrazinyl.

131. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 7, wherein

R^3 represents unsubstituted or singly or multiply substituted thiophenyl, benzothiophenyl, pyridyl, furyl, benzofuranyl, benzodioxolanyl, indolyl, indanyl, benzodioxanyl, pyrrolyl, pyrimidyl or pyrazinyl.

132. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 5, wherein R^3 represents phenyl, naphthyl or anthracenyl.

133. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 6, wherein R^3 represents phenyl, naphthyl or anthracenyl.

134. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 1, wherein

R⁴ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl; or -R⁸-L-R⁹.

135. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 134, wherein

R⁸ represents unsubstituted or singly or multiply substituted indolyl, naphthyl, benzofuranyl, benzothiophenyl, indanyl, benzodioxanyl, benzodioxolanyl, acenaphthyl, carbazolyl, phenyl, thiophenyl, furyl, pyridyl, pyrrolyl, pyrazinyl or pyrimidyl, fluorenlyl, fluoranthenyl, benzothiazolyl, benzotriazolyl or benzo[1,2,5]thiazolyl or 1,2-dihydroacenaphthenyl, pyridinyl, furanyl, benzofuranyl, pyrazolinonyl, oxopyrazolinonyl, pyrimidinyl, quinolinyl, isoquinolinyl, phthalazinyl or quinazolinyl;

L represents -C(O)-NH-, -NH-C(O)-, -C(O)-O-, -O-C(O)-, -O-, -S-or -S(O)₂-;

or R⁹ represents unsubstituted or singly or multiply substituted indolyl, naphthyl, benzofuranyl, benzothiophenyl, indanyl, benzodioxanyl, benzodioxolanyl, acenaphthyl, carbazolyl, phenyl, thiophenyl, furyl, pyridyl, pyrrolyl, pyrazinyl or pyrimidyl, fluorenlyl, fluoranthenyl, benzothiazolyl, benzotriazolyl or benzo[1,2,5]thiazolyl or 1,2-dihydroacenaphthenyl, pyridinyl, furanyl, benzofuranyl, pyrazolinonyl, oxopyrazolinonyl, pyrimidinyl, quinolinyl, isoquinolinyl, phthalazinyl or quinazolinyl.

136. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 1, wherein

R⁴ represents -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷.

137. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 136, wherein

R⁶ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₄-alkyl; or saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₄-alkyl.

138. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 136, wherein

R⁷ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl.

139. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 8, wherein

R⁴ represents unsubstituted or singly or multiply substituted heteroaryl; or -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

where Y = H₂.

140. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 139, wherein

R⁶ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₄-alkyl;

or

R⁷ represents unsubstituted or singly or multiply substituted indolyl, benzofuranyl, benzothiophenyl, indanyl, benzodioxanyl, benzodioxolanyl, acenaphthyl, carbazolyl, thiophenyl, furyl, pyridyl, pyrrolyl, pyrazinyl or pyrimidyl, fluorenyl, fluoranthenyl, benzothiazolyl, benzotriazolyl or benzo[1,2,5]thiazolyl or 1,2-dihydroacenaphthenyl, pyridinyl, furanyl,

benzofuranyl, pyrazolinonyl, oxopyrazolinonyl, pyrimidinyl, quinolinyl, isoquinolinyl, phthalazinyl or quinazolinyl.

141. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 139, wherein

R⁸ represents unsubstituted or singly or multiply substituted indolyl, naphthyl, benzofuranyl, benzothiophenyl, indanyl, benzodioxanyl, benzodioxolanyl, acenaphthyl, carbazolyl, phenyl, thiophenyl, furyl, pyridyl, pyrrolyl, pyrazinyl or pyrimidyl, fluorenlyl, fluoranthenyl, benzothiazolyl, benzotriazolyl or benzo[1,2,5]thiazolyl or 1,2-dihydroacenaphthenyl, pyridinyl, furanyl, benzofuranyl, pyrazolinonyl, oxopyrazolinonyl, pyrimidinyl, quinolinyl, isoquinolinyl, phthalazinyl or quinazolinyl;

or R⁹ represents unsubstituted or singly or multiply substituted indolyl, naphthyl, benzofuranyl, benzothiophenyl, indanyl, benzodioxanyl, benzodioxolanyl, acenaphthyl, carbazolyl, phenyl, thiophenyl, furyl, pyridyl, pyrrolyl, pyrazinyl or pyrimidyl, fluorenlyl, fluoranthenyl, benzothiazolyl, benzotriazolyl or benzo[1,2,5]thiazolyl or 1,2-dihydroacenaphthenyl, pyridinyl, furanyl, benzofuranyl, pyrazolinonyl, oxopyrazolinonyl, pyrimidinyl, quinolinyl, isoquinolinyl, phthalazinyl or quinazolinyl.

142. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 9, wherein

R⁴ represents -CHR⁶R⁷, -CHR⁶-CH₂R⁷ or -CHR⁶-CH₂-CH₂R⁷.

143. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 142, wherein

R⁶ represents saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₄-alkyl.

or

R⁷ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl.

144. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 10, wherein

R⁴ represents -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷

where Y = O.

145. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 144, wherein

R⁷ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl.

146. (Previously Presented) A substituted 4-aminocyclohexanol compound according to claim 1, wherein said compound is selected from the group consisting of:

- 4-dimethylamino-1-(1-methyl-1H-indol-2-yl)-4-phenylcyclohexanol
- 1-benzo[b]thiophen-2-yl-4-dimethylamino-4-phenylcyclohexanol
- 1-benzo[b]thiophen-3-yl-4-dimethylamino-4-phenylcyclohexanol
- 1-(1-benzenesulphonyl-1H-indol-2-yl)-4-dimethylamino-4-phenylcyclohexanol
- 1-benzofuran-2-yl-4-dimethylamino-4-phenylcyclohexanol; and
- 1-benzothiazol-2-yl-4-dimethylamino-4-phenylcyclohexanol, or


a salt thereof with a physiologically tolerated acid.

147. (Previously Presented) A pharmaceutical composition containing at least one substituted 4-aminocyclohexanol compound according to claim 1.

148. (Previously Presented) A pharmaceutical composition according to claim 147, wherein the pharmaceutical composition comprises an opioid or an anesthetic.

149. (Previously Presented) A method of alleviating pain or treating a locomotive disorder or administering an anticonvulsant or muscle relaxant, said method comprising the step of administering to a mammal in need thereof an effective amount of a compound according to claim 1.

150. (Previously Presented) A method of treating phobias, stress and syndromes associated with stress, depression, epilepsy, Alzheimer's disease, senile dementia, general cognitive dysfunction, learning difficulties and memory loss, withdrawal symptoms, alcohol abuse or dependency, drug abuse or dependency, sexual dysfunction, cardiovascular diseases, hypotension, hypertension, tinnitus, pruritus, defective hearing, defective bowel motility, impaired assimilation of food, anorexia, obesity, diarrhoea, cachexia, urinary incontinence or for providing an antitussive or anaesthetic or for co-administration during treatment with an opioid analgesic or with an anaesthetic, for diuresis, antinatriuresis or anxiolysis in a mammal, said method comprising administering to said mammal an effective amount of a substituted 4-aminocyclohexanol compound corresponding to formula I,



I

wherein

R¹ and R² independently of one another represent H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl; wherein R¹ and R² are not both H, or the radicals R¹ and R² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁵CH₂CH₂ or (CH₂)₃₋₆;

wherein R⁵ represents H; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted alkylene aryl bound via C₁₋₃, C₃₋₈-cycloalkyl or heteroaryl;

R³ represents unsubstituted or singly or multiply substituted aryl or heteroaryl; R⁴ represents unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl; -CHR⁶R⁷, -CHR⁶-CH₂R⁷, -CHR⁶-CH₂-CH₂R⁷, -CHR⁶-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

wherein

Y = O, S or H₂;

R⁶ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₇-alkyl; or saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C(O)O-C₁₋₆-alkyl;

R⁷ represents H; respectively unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl;

R⁸ represents respectively unsubstituted or singly or multiply substituted aryl or heteroaryl;


L represents -C(O)-NH-, -NH-C(O)-, -C(O)-O-, -O-C(O)-, -O-, -S- or -S(O)₂-; and

R^9 represents unsubstituted or singly or multiply substituted aryl or heteroaryl,


or a salt thereof with a physiologically tolerated acid.

151. (Previously Presented) A method of producing a substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1 comprising the steps of:


a. reacting a cyclohexane-1,4-dione protected by the groups S¹ and S² according to formula II in the presence of a compound corresponding to formula HNR⁰¹R⁰² with a cyanide, to form a protected N-substituted 1-amino-4-oxo-cyclohexanecarbonitrile compound corresponding to formula III;




b. reacting the compound corresponding to formula III with organometallic reagents corresponding to the formula metal-R³ to form a compound corresponding to formula IVa;



c. removing the protective groups S^1 and S^2 on the compound corresponding to formula IVa to form a 4-substituted 4-aminocyclohexanone compound corresponding to formula IV;



d. reacting the 4-substituted 4-aminocyclohexanone compound corresponding to formula IV with organometallic reagents corresponding to the formula metal-R⁰⁴ to form a compound corresponding to formula V;



wherein

R⁰¹ and R⁰² independently of one another represent H; H provided with a protective group; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or

multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl;

or the radicals R⁰¹ and R⁰² together form a ring and represent CH₂CH₂OCH₂CH₂, CH₂CH₂NR⁰⁵CH₂CH₂ or (CH₂)₃₋₆,

wherein R⁰⁵ represents H; H provided with a protective group; saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₈-alkyl or C₃₋₈-cycloalkyl; singly or multiply substituted or unsubstituted aryl or heteroaryl; or singly or multiply substituted or unsubstituted aryl bound via C₁₋₃-alkylene, C₃₋₈-cycloalkyl or heteroaryl;

R⁰⁴ represents H, H provided with a protective group; unsubstituted or singly or multiply substituted C₃₋₈-cycloalkyl, aryl or heteroaryl; -CHR¹⁰R⁷, -CHR¹⁰-CH₂R⁷, -CHR¹⁰-CH₂-CH₂R⁷, -CHR¹⁰-CH₂-CH₂-CH₂R⁷, -C(Y)R⁷, -C(Y)-CH₂R⁷, -C(Y)-CH₂-CH₂R⁷ or -C(Y)-CH₂-CH₂-CH₂R⁷; or -R⁸-L-R⁹

where R¹⁰ represents H, saturated or unsaturated, branched or unbranched, singly or multiply substituted or unsubstituted C₁₋₇-alkyl; and

S¹ and S² independently of one another represent protective groups or together represent a protective group.

152. (Previously Presented) The method of claim 151, wherein S¹ and S² together represent a monoacetal group.

153. (currently amended) The method of claim 151, wherein step a) further comprises:

acylating, alkylating or sulfonating the compound corresponding to formula III in any sequence and optionally repeatedly; or

where R^{01} or R^{02} or $[[R^{06}]] \underline{R^6} = H$ protected with a protective group, removing at least one protective group and optionally acylating, alkylating or sulfonating the compound corresponding to formula III; or

where R^{01} or R^{02} or $[[R^{06}]] \underline{R^6} = H$, introducing at least one protective group and optionally acylating, alkylating or sulfonating the compound corresponding to formula III.

154. (currently amended) The method of claim 151, wherein step b) further comprises:

acylating, alkylating or sulfonating the compound corresponding to formula IVa in any sequence and optionally repeatedly; or

where R^{01} , R^{02} or $[[R^{06}]] \underline{R^6} = H$ protected with a protective group, removing at least one protective group and optionally acylating, alkylating or sulfonating the compound corresponding to formula IVa; or

where R^{01} or R^{02} or $[[R^{06}]] \underline{R^6} = H$, introducing at least one protective group and optionally acylating, alkylating or sulfonating the compound corresponding to formula IVa.

155. (currently amended) The method of claim 151, wherein step c) further comprises:

acylating, alkylating or sulfonating the compound corresponding to formula IV in any sequence and optionally repeatedly; or

where R^{01} , R^{02} or $[[R^{06}]] \underline{R^6} = H$ protected with a protective group, removing at least one protective group and optionally acylating, alkylating or sulfonating the compound corresponding to formula IV; or

where R^{01} or R^{02} or $[[R^{06}]] \underline{R^6} = H$, introducing at least one protective group and optionally acylating, alkylating or sulfonating the compound corresponding to formula IV.

156. (currently amended) The method of claim 151, wherein step d) further comprises:

acylating, alkylating or sulfonating the compound corresponding to formula V in any sequence and optionally repeatedly; or

where R^{01} , R^{02} or $[[R^{06}]]$ R^6 = H protected with a protective group, removing at least one protective group and optionally acylating, alkylating or sulfonating the compound corresponding to formula V; or

where R^{01} or R^{02} or $[[R^{06}]]$ R^6 = H, introducing at least one protective group and optionally acylating, alkylating or sulfonating the compound corresponding to formula V.

157. (Previously Presented) The method of claim 151, wherein the protective groups on H in R^{01} , R^{02} , R^{04} or R^{05} are selected from the group consisting of alkyl groups, benzyl groups or carbamates.

158. (Previously Presented) The method of claim 157, wherein the protective groups on H in R^{01} , R^{02} , R^{04} or R^{05} are selected from the group consisting of fluorenylmethyl-chloroformate groups (FMOC), benzyloxycarbonyl (Z) and tert-butyloxycarbonyl (Boc).

159. (Previously Presented) The method of claim 151, wherein the cyanide of step a) is potassium cyanide.

160. (Previously Presented) The method of claim 151, wherein the organometallic reagents of step b) are Grignard or organolithium reagents.

161. (Previously Presented) The method of claim 151, wherein the organometallic reagents of step d) are Grignard or organolithium reagents.

162. (New) The substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1, wherein said compound is in the form of a salt with a physiologically tolerated acid.

163. (New) The substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1, wherein said compound is in the form of a hydrate.

164. (New) The substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1, wherein said compound is present in the form of a pure enantiomer or pure diastereoisomer.

165. (New) The substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1, wherein said compound is present in the form of a mixture of stereoisomers.

166. (New) The substituted 4-aminocyclohexanol compound corresponding to formula I of claim 1, wherein said compound is present in the form of a racemic mixture.